

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Recombinant Cynomolgus IL-31 Protein, Human ABB-RP02727

Recombinant Cynomolgus IL-31 Protein, Human
ABB-RP02727
RP02727
ABB-RP02727-100UG
ABclonal
Human
Proteine/Peptide
Monkey
Leu27-Thr163
Many autoimmune skin diseases, such as bullous pemphigoid (BP), psoriasis and certain types of chronic urticaria, are associated with intensive pruritus. While histamine and neuropeptides have previously been ascribed to play a role in itch that acco
$< 1$ EU/ $\mu$ g of the protein by LAL method.
16.5 kDa
102135935
G7PJ78
95 % as determined by Tris-Bis PAGE, 95 % as determined by HPLC.
Cynomolgus IL-31

**Application Notes** 

Cross-Reactivity: Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA,5% HSA,10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles. ResearchArea: Other Recombinant Protein