

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Human PDCD1 protein BYT-ORB705336

|                            |   |
|----------------------------|---|
| Article Name               | Human PDCD1 protein   |
| Biozol Catalog Number      | BYT-ORB705336   |
| Supplier Catalog Number    | orb705336   |
| Alternative Catalog Number | BYT-ORB705336-1,BYT-ORB705336-100,BYT-ORB705336-20  |
| Manufacturer               | Biorbyt   |
| Category                   | Proteine/Peptide  |
| Product Description        | This Human PDCD1 protein spans the amino acid sequence from region 25-167aa. Purity: Greater than 93% as determined by SDS-PAGE....                   |
| Molecular Weight           | 18.2 kDa  |
| UniProt                    | <a href="#">Q15116</a>  |
| Buffer                     | Lyophilized from a 0.2 µm filtered 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0   |
| Source                     | Homo sapiens (Human)  |
| Purity                     | Greater than 93% as determined by SDS-PAGE.   |
| Form                       | Lyophilized powder  |
| Sequence                   | LDSPDRPWNPPTFSPALLVVTEGDNATFTCSFSNTSESVLNWYRMSPSNQT<br>DKLAAPFEDRSQPGQDCRFRVTQLPNGRDFHMSVVRARRNDSGYLCGAISL<br>APKAQIKESLRAELRVTERRAEVPTAHPSPSPRPAGQFQ |

|                   |   |
|-------------------|---|
| Application Notes | <p>Biological Origin: Homo sapiens (Human). Biological Activity: Measured by its binding ability in a functional ELISA. Immobilized PD-1 at 2 µg/ml can bind Anti-PD-1 recombinant antibody, the EC50 of human PD-1 protein is 6.087-7.854 ng/ml. ②Measured by its binding ability in a functional ELISA. Immobilized PD-1 at 2 µg/ml can bind Nivolumab, the EC50 of human PD-1 protein is 9.713-12.39 ng/ml.</p> <p>Application Notes: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference</p> |
|-------------------|---|