

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD155 Antibody, Clone: [SKII.4], Unconjugated, Mouse, Monoclonal BYT-ORB2988638

| Article Name | CD155 Antibody, Clone: [SKII.4], Unconjugated, Mouse, Monoclonal |
|----------------------------|---|
| Biozol Catalog Number | BYT-ORB2988638 |
| Supplier Catalog Number | orb2988638 |
| Alternative Catalog Number | BYT-ORB2988638-30,BYT-ORB2988638-50,BYT-ORB2988638- 100,BYT-ORB2988638-200 |
| Manufacturer | Biorbyt |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF |
| Species Reactivity | Human |
| Immunogen | SK-N-S1 human neuroblastoma cells |
| Conjugation | Unconjugated |
| Product Description | The CD155 Antibody is suitable for FC, IF. It is a Monoclonal, Unconjugated antibody which raised against SK-N-S1 human neuroblastoma cells .Purification: The antibody was purified by affinity chromatography |
| Clonality | Monoclonal |
| Clone Designation | [SKII.4] |
| UniProt | P15151 |
| Form | Mouse IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide. |

Application Notes

Application Notes: 1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube. 2. Add appropriate amount of antibody to the bottom of flow tube mixing with the whole blood, incubate for 30 minutes at room temperature. 3. Add 2 ml RBC lysis buffer, incubate for 10 minutes after mixing, dissolve red blood cells. 4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant. 5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant. 6. Add appropriate amount of fluorescentlabeled anti-mouse IgGs and incubate for 20 minutes away from light at room temperature. 7. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant. 8. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 C then measured)