

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Recombinant Mouse CCL20/MIP-3-alpha Protein, E. coli ABB-RP03272

|                            |   |
|----------------------------|---|
| Article Name               | Recombinant Mouse CCL20/MIP-3-alpha Protein, E. coli  |
| Biozol Catalog Number      | ABB-RP03272   |
| Supplier Catalog Number    | RP03272   |
| Alternative Catalog Number | ABB-RP03272-20UG, ABB-RP03272-100UG   |
| Manufacturer               | ABclonal  |
| Host                       | E. coli   |
| Category                   | Proteine/Peptide  |
| Species Reactivity         | Mouse   |
| Immunogen                  | Ala28-Met97   |
| Product Description        | Chemokine (C-C motif) ligand 2 (CCL2) or liver activation regulated chemokine (LARC) or Macrophage Inflammatory Protein-3 (MIP3A) is a small cytokine belonging to the CC chemokine family that attracts immature dendritic cells and memory T lymphocytes... |
| Concentration              | Please contact us for more information.   |
| Molecular Weight           | 8.1 kDa   |
| NCBI                       | <a href="#">20297</a>   |
| UniProt                    | <a href="#">O89093</a>  |
| Purity                     | 95 % as determined by SDS-PAGE.   |
| Form                       | Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.   |

|                    |   |
|--------------------|---|
| Target             | CCL20   |
| Application Dilute | Lyophilized from a 0.22 $\mu$ m filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.  |
| Application Notes  | Cross-Reactivity: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles., ResearchArea: Cytokines & Cytokine Receptors |