

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human S100 beta protein, His tag, Unconjugated GTX00117-PRO

Article Name	Human S100 beta protein, His tag, Unconjugated
Biozol Catalog Number	GTX00117-PRO
Supplier Catalog Number	GTX00117-pro
Alternative Catalog Number	GTX00117-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	6285
UniProt	P04271
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	Full length protein, N-terminal His-Tag, Met1~Glu92 (NP_006263.1)

Application Notes

Protein S100B is a member of the S100 family. S100 proteins are EF-hand calcium-binding proteins and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. Experimental results suggest that the receptor for advanced glycation end products (RAGE) plays important roles in mediating S100 protein-induced cellular signaling. Thus a binding ELISA assay was conducted to detect the interaction of recombinant human S100B and recombinant human RAGE. Briefly, S100B were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to RAGE-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-S100B mAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of of S100B and RAGE was in a dose dependent manner.