

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Human CD137 protein, His tag, Unconjugated GTX00128-PRO

Article Name	Human CD137 protein, His tag, Unconjugated
Biozol Catalog Number	GTX00128-PRO
Supplier Catalog Number	GTX00128-pro
Alternative Catalog Number	GTX00128-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	<a href="#">3604</a>
UniProt	<a href="#">Q07011</a>
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Asn40~Glu153 (NP_001552.2)

Application Notes

TNFRSF9 (Tumor necrosis factor receptor superfamily member 9) is a member of the TNF-receptor superfamily. This receptor contributes to the clonal expansion, survival, and development of T cells. A binding ELISA assay was conducted to detect the interaction of TNFRSF9 and TNFa. Briefly, recombinant human TNFRSF9 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100  $\mu$ l were then transferred to TNFa-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-TNFRSF9 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50  $\mu$ l stop solution to the wells and read at 450nm immediately. The binding activity of TNFRSF9 and TNFa was in a dose dependent manner.