

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human Arylsulfatase F protein, His tag, Unconjugated GTX00162-PRO

Article Name	Human Arylsulfatase F protein, His tag, Unconjugated
Biozol Catalog Number	GTX00162-PRO
Supplier Catalog Number	GTX00162-pro
Alternative Catalog Number	GTX00162-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	416
UniProt	P54793
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	HEK293 cells
Form	Lyophilized powder
Sequence	N-terminal His-Tag, His23~Asp300 (NP_001188467.1)

Application Notes

ARSF (Arylsulfatase F) is a member of the sulfatase gene family, and more specifically, the arylsulfatase subfamily. Sulfatases are essential for the correct composition of bone and cartilage matrix. The activity of this protein, unlike that of arylsulfatase E, is not inhibited by warfarin. Besides, EP300 (E1A binding protein p300) has been identified as an interactor of ARSF, thus a binding ELISA assay was conducted to detect the interaction of recombinant human ARSF and recombinant human EP300. Briefly, ARSF were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to EP300-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-ARSF pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of ARSF and EP300 was in a dose dependent manner.