

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human Leptin protein, His and GST tag, Unconjugated GTX00170-PRO

Article Name	Human Leptin protein, His and GST tag, Unconjugated
Biozol Catalog Number	GTX00170-PRO
Supplier Catalog Number	GTX00170-pro
Alternative Catalog Number	GTX00170-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	3952
UniProt	P41159
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His and GST-Tag, Val22~Cys167 (NP_000221.1)

Application Notes

Leptin is a hormone made by adipose cells that helps to regulate energy balance by inhibiting hunger. Many of leptin's effects are mediated through neuropeptide-containing neurons and neuropeptide receptors in the hypothalamus. Although regulation of fat stores is deemed to be the primary function of leptin, it also plays a role in other physiological processes, as evidenced by its multiple sites of synthesis other than fat cells, and the multiple cell types beside hypothalamic cells that have leptin receptors. Besides, Hexokinase 3, White Cell (HK3) has been identified as an interactor of LEP, thus a binding ELISA assay was conducted to detect the interaction of recombinant human LEP and recombinant human HK3. Briefly, LEP were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to HK3-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-LEPpAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of LEP and HK3 was in a dose dependent manner.