

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human MDM2 protein, His tag, Unconjugated GTX00176-PRO

Article Name	Human MDM2 protein, His tag, Unconjugated
Biozol Catalog Number	GTX00176-PRO
Supplier Catalog Number	GTX00176-pro
Alternative Catalog Number	GTX00176-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	4193
UniProt	Q00987
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Met1~Pro218 (NP_001138811.1)

Application Notes

Mouse double minute 2 homolog (MDM2) also known as E3 ubiquitin-protein ligase Mdm2 is a cellular oncoprotein that recognizes the N-terminal trans-activation domain (TAD) of the p53 tumor suppressor and as an inhibitor of p53 transcriptional activation. The human homologue of this protein is sometimes called Hdm2. The p53 tumor suppressor is the key target of MDM2. It has been identified as a p53 interacting protein that represses p53 transcriptional activity and also acts as an E3 ubiquitin ligase, targeting both itself and p53 for degradation by the proteasome. MDM2 is capable of auto-polyubiquitination, and in complex with p300, a cooperating E3 ubiquitin ligase, is capable of polyubiquitinating p53. Besides, S100 Calcium Binding Protein (S100) has been identified as an interactor of MDM2, thus a binding ELISA assay was conducted to detect the interaction of recombinant human MDM2 and recombinant human S100. Briefly, MDM2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to S100-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-MDM2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of MDM2 and S100 was in a dose dependent manner.