

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human IL29 protein, His tag, Unconjugated GTX00245-PRO

Article Name	Human IL29 protein, His tag, Unconjugated
Biozol Catalog Number	GTX00245-PRO
Supplier Catalog Number	GTX00245-pro
Alternative Catalog Number	GTX00245-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	282618
UniProt	Q8IU54
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Gly20~Thr200 (NP_742152.1)

Application Notes

Interleukin-29 (IL-29) is a member of the helical cytokine family and is a type III interferon. It is also known as IFN λ 1 and is highly similar in amino acid sequence to the IL-28, the other type III interferon. IL-29 plays an important role in host defenses against microbes and its gene is highly upregulated in cells infected with viruses. Besides, ATPase, Ca Transporting, Plasma Membrane 2 (ATP2B2) has been identified as an interactor of IL29, thus a binding ELISA assay was conducted to detect the interaction of recombinant human IL29 and recombinant human ATP2B2. Briefly, IL29 were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to ATP2B2-coated microtiter wells and incubated for 2h at 37 C. Wells were washed with PBST and incubated for 1h with anti-IL29 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of IL29 and ATP2B2 was in a dose dependent manner.