

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse Arginase 1 protein, His tag, Unconjugated GTX00303-PRO

Article Name	Mouse Arginase 1 protein, His tag, Unconjugated
Biozol Catalog Number	GTX00303-PRO
Supplier Catalog Number	GTX00303-pro
Alternative Catalog Number	GTX00303-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Mouse
Conjugation	Unconjugated
NCBI	11846
UniProt	Q61176
Buffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	Full length protein, N-terminal His-Tag, Met1~Lys323 (NP_031508.1)

Application Notes

Arginase (Arg) is an enzyme that catalyzes the degradation of arginine to produce urea and ornithine, which is crucial in the urea cycle. In most mammals, two isozymes of this enzyme exist, the first, Arginase I, functions in the urea cycle, and is located primarily in the cytoplasm of the liver. The second isozyme, Arginase II, has been implicated in the regulation of the arginine/ornithine concentrations in the cell. Besides, Ubiquitin Carboxyl Terminal Hydrolase L5 (UCHL5) has been identified as an interactor of Arg, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse Arg and recombinant mouse UCHL5. Briefly, Arg were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to UCHL5-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-Arg pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of Arg and UCHL5 was in a dose dependent manner.