

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human MIP3 beta protein, His tag (active), Unconjugated GTX00424-PRO

Article Name	Human MIP3 beta protein, His tag (active), Unconjugated
Biozol Catalog Number	GTX00424-PRO
Supplier Catalog Number	GTX00424-pro
Alternative Catalog Number	GTX00424-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA
Species Reactivity	Human
Conjugation	Unconjugated
NCBI	6363
UniProt	Q99731
Buffer	Reconstitute with 20mM Tris (pH8.0) and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris (pH8.0), 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Form	Lyophilized powder
Sequence	N-terminal His-Tag, Gly22~Ser98 (NP_006265.1)

Application Notes

Macrophage Inflammatory Protein 3 Beta (MIP3b) is a small cytokine belonging to the CC chemokine family that is also known as EBI1 ligand chemokine (ELC) and Chemokine C-C motif ligand 19 (CCL19). This chemokine elicits its effects on its target cells by binding to the chemokine receptor chemokine receptor CCR7. It attracts certain cells of the immune system, including dendritic cells and antigen-engaged B cells, CCR7 central-memory T-Cells. Thus, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of recombinant human MIP3b on the Jurkat cell line. Briefly, Jurkat cells were seeded into the upper chambers (150 μ l cell suspension, 106 cells/ml in RPMI-1640 with FBS free) and MIP3b (0.01, 0.1, 1, 10, 100, and 1000 ng/ml diluted separately in serum free RPMI-1640) was added in lower chamber with a polycarbonate filter (8 μ m pore size) used to separate the two compartments. After incubation at 37C with 5% CO₂ for 3h, the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification (*100) and the number of migrated cells were counted at high magnification (*400) randomly (five fields for each filter). MIP3b is able to induce migration of Jurkat cells, and the migrated cells were counted at high magnification (*400). The optimum chemotaxis of recombinant human MIP3b occurs at 0.1-1 ng/ml.