

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse Annexin II protein, His and TrxA tag, Unconjugated GTX03782-PRO

Article Name	Mouse Annexin II protein, His and TrxA tag, Unconjugated
Biozol Catalog Number	GTX03782-PRO
Supplier Catalog Number	GTX03782-pro
Alternative Catalog Number	GTX03782-PRO-10
Manufacturer	GeneTex
Category	Proteine/Peptide
Application	FA, WB
Species Reactivity	Mouse
Conjugation	Unconjugated
NCBI	12306
UniProt	P07356
Buffer	Reconstitute with 10mM PBS (pH7.4) to 0.1-1.0 mg/mL. Do not vortex. Lyophilized from PBS pH7.4, 0.01% SKL, 5% Trehalose, no preservatives.
Expression System	E. coli
Form	Lyophilized powder
Sequence	Full length protein, N-terminal His and TrxA Tag, Met1~Asp339 (NP_031611.1)

Application Notes

Annexin A2 (ANXA2) also known as annexin II is a member of the annexin family. Members of this calcium-dependent phospholipid-binding protein family play a role in the regulation of cellular growth and in signal transduction pathways. This protein functions as an autocrine factor which heightens osteoclast formation and bone resorption. It is a calcium-dependent phospholipid-binding protein whose function is to help organize exocytosis of intracellular proteins to the extracellular domain. Besides, S100 Calcium Binding Protein A10 (S100A10) has been identified as an interactor of ANXA2, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse ANXA2 and recombinant mouse S100A10. Briefly, ANXA2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to S100A10-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-ANXA2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of ANXA2 and S100A10 was in a dose dependent manner.