

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### CD8a Antibody, Clone: [SPM548], Mouse, Monoclonal NSJ-V2386IHC-7ML

Article Name	CD8a Antibody, Clone: [SPM548], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2386IHC-7ML
Supplier Catalog Number	V2386IHC-7ML
Alternative Catalog Number	NSJ-V2386IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Human CD8 recombinant protein was used as immunogen for this CD8a antibody.
Product Description	CD8 molecule consists of two chains, termed alpha and beta chain, which are expressed as a disulphide-linked heterodimer or homodimer. CD8 is expressed on T cell subset (cytotoxic/suppressor T cells), thymocytes and NK cells....
Clonality	Monoclonal
Clone Designation	[SPM548]
UniProt	<a href="#">P01732</a>
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Variations in protocols, secondaries and substrates may require the CD8a antibody to be titrated up or down for optimal performance.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.