

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CD20 Antibody, Clone: [SPM618], Mouse, Monoclonal NSJ-V2390IHC-7ML

Article Name	Anti-CD20 Antibody, Clone: [SPM618], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2390IHC-7ML
Supplier Catalog Number	V2390IHC-7ML
Alternative Catalog Number	NSJ-V2390IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human MS4A1/CD20 protein was used as the immunogen for this anti-CD20 antibody.
Product Description	This antibody recognizes a protein of 33-37kDa, identified as CD20. The epitope is similar to or identical to that recognized by other CD20 antibodies including Leu-16 and B1. This antibody can be used for immunophenotyping of leukemia and malignant ...
Clonality	Monoclonal
Clone Designation	[SPM618]
UniProt	P11836
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the anti-CD20 antibody to be titered up or down for optimal performance.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.