

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

GCSF Antibody, Clone: [SPM468], Mouse, Monoclonal NSJ-V2472-100UG

Article Name	GCSF Antibody, Clone: [SPM468], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2472-100UG
Supplier Catalog Number	V2472-100UG
Alternative Catalog Number	NSJ-V2472-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Nuclei from pokeweed mitogen stimulated human peripheral blood lymphocytes was used as the immunogen for the GCSF antibody.
Product Description	This mAb recognizes granulocyte-colony stimulating factor (G-CSF) in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell types. Markers of myeloid cells are useful in the identification of different levels of cellular dif...
Clonality	Monoclonal
Clone Designation	[SPM468]
UniProt	P09919
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	<p>Optimal dilution of the GCSF antibody should be determined by the researcher.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>