

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Secretory Component Glycoprotein Antibody / ECM1, Clone: [rECM1/792], Mouse, Monoclonal NSJ-V2484SAF-100UG

| | |
|----------------------------|---|
| Article Name | Recombinant Secretory Component Glycoprotein Antibody / ECM1, Clone: [rECM1/792], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V2484SAF-100UG |
| Supplier Catalog Number | V2484SAF-100UG |
| Alternative Catalog Number | NSJ-V2484SAF-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC-P |
| Species Reactivity | Human |
| Immunogen | Recombinant human protein was used as the immunogen for the recombinant Secretory Component Glycoprotein antibody. |
| Product Description | This mAb reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free... |
| Clonality | Monoclonal |
| Clone Designation | [rECM1/792] |
| UniProt | Q16610 |
| Purity | Protein G affinity chromatography |

| | |
|--------------------|---|
| Form | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1),Prediluted IHC only format: incubate for 30 min at RT (2) |
| Application Notes | <p>Optimal dilution of the recombinant Secretory Component Glycoprotein antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |