

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CD99 Antibody, Clone: [SPM596], Mouse, Monoclonal NSJ-V2710-20UG

| | |
|----------------------------|---|
| Article Name | Anti-CD99 Antibody, Clone: [SPM596], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V2710-20UG |
| Supplier Catalog Number | V2710-20UG |
| Alternative Catalog Number | NSJ-V2710-20UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC-P |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human protein was used as the immunogen for the anti-CD99 antibody. |
| Product Description | Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which are produced by a... |
| Clonality | Monoclonal |
| Clone Designation | [SPM596] |
| UniProt | P14209 |
| Purity | Protein G affinity chromatography |

| | |
|--------------------|---|
| Form | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Application Notes | <p>Optimal dilution of the anti-CD99 antibody should be determined by the researcher.</p> <ol style="list-style-type: none"> 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |