

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CD99 Antibody, Clone: [SPM596], Mouse, Monoclonal NSJ-V2710SAF-100UG

Article Name	Anti-CD99 Antibody, Clone: [SPM596], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2710SAF-100UG
Supplier Catalog Number	V2710SAF-100UG
Alternative Catalog Number	NSJ-V2710SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the anti-CD99 antibody.
Product Description	Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which are produced by a...
Clonality	Monoclonal
Clone Designation	[SPM596]
UniProt	P14209
Purity	Protein G affinity chromatography
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free

Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	<p>Optimal dilution of the anti-CD99 antibody should be determined by the researcher.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>