

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CDw17 Antibody / Lactosylceramide / LaCer, Clone: [HuLy-m13], Mouse, Monoclonal NSJ-V3052SAF-100UG

| | |
|----------------------------|---|
| Article Name | CDw17 Antibody / Lactosylceramide / LaCer, Clone: [HuLy-m13], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V3052SAF-100UG |
| Supplier Catalog Number | V3052SAF-100UG |
| Alternative Catalog Number | NSJ-V3052SAF-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF |
| Species Reactivity | Human |
| Immunogen | Beta-2 Microglobulin associated proteins from a detergent lysate of human PBLs were used as the immunogen for the CDw17 antibody. |
| Product Description | CD17/CDw17 is an intermediate glycosphingolipid from the metabolism of higher gangliosides that localizes to sphingolipid-sterol rafts. CD17 is detectable in monocytes, granulocytes, basophils, platelets, a subset of peripheral B cells (CD19+) and to... |
| Clonality | Monoclonal |
| Clone Designation | [HuLy-m13] |
| UniProt | Not known |
| Purity | PEG precipitation |

| | |
|--------------------|--|
| Form | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Dilute | Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 0.5-1ug/ml |
| Application Notes | Optimal dilution of the CDw17 antibody should be determined by the researcher. |