

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 10/13 Antibody, Clone: [DE-K13], Mouse, Monoclonal NSJ-V3053-100UG

| | |
|----------------------------|---|
| Article Name | Cytokeratin 10/13 Antibody, Clone: [DE-K13], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V3053-100UG |
| Supplier Catalog Number | V3053-100UG |
| Alternative Catalog Number | NSJ-V3053-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF, IHC-P, WB |
| Species Reactivity | Human |
| Immunogen | A cytoskeletal preparation extracted from human ectocervical epithelium was used as the immunogen for the Cytokeratin 10/13 antibody. |
| Product Description | This antibody recognizes cytokeratin 10 (56.5kDa) and cytokeratin 13 (53kDa) in Western blotting. It recognizes only cytokeratin 13 in formalin-fixed, paraffin-embedded tissue sections. It does not react with cytokeratin 10 positive, cytokeratin 13 n... |
| Clonality | Monoclonal |
| Clone Designation | [DE-K13] |
| UniProt | P13645 |
| Purity | Protein G affinity chromatography |

| | |
|--------------------|--|
| Form | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Dilute | Flow cytometry: 1-2ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Application Notes | Optimal dilution of the Cytokeratin 10/13 antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |