

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 10/13 Antibody, Clone: [DE-K13], Mouse, Monoclonal NSJ-V3053IHC-7ML

Article Name	Cytokeratin 10/13 Antibody, Clone: [DE-K13], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V3053IHC-7ML
Supplier Catalog Number	V3053IHC-7ML
Alternative Catalog Number	NSJ-V3053IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	A cytoskeletal preparation extracted from human ectocervical epithelium was used as the immunogen for the Cytokeratin 10/13 antibody.
Product Description	This antibody recognizes cytokeratin 10 (56.5kDa) and cytokeratin 13 (53kDa) in Western blotting. It recognizes only cytokeratin 13 in formalin-fixed, paraffin-embedded tissue sections. It does not react with cytokeratin 10 positive, cytokeratin 13 n...
Clonality	Monoclonal
Clone Designation	[DE-K13]
UniProt	P13645
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the Cytokeratin 10/13 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.