

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **hCG Holo Antibody, Clone: [HCGab/52], Mouse, Monoclonal NSJ-V3064-100UG**

|                            |   |
|----------------------------|---|
| Article Name               | hCG Holo Antibody, Clone: [HCGab/52], Mouse, Monoclonal   |
| Biozol Catalog Number      | NSJ-V3064-100UG   |
| Supplier Catalog Number    | V3064-100UG   |
| Alternative Catalog Number | NSJ-V3064-100UG   |
| Manufacturer               | NSJ Bioreagents   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC-P   |
| Species Reactivity         | Human   |
| Immunogen                  | Purified hCG protein was used as the immunogen for the hCG Holo antibody.   |
| Product Description        | This mAb reacts only with the intact-HCG and not with either free alpha- or free beta-chain of HCG. HCG is a glycoprotein and is composed of two non-identical, non-covalently linked polypeptide chains designated as the alpha and beta subunits. The al... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [HCGab/52]  |
| UniProt                    | <a href="#">P01215</a>  |
| Purity                     | Protein G affinity chromatography   |

|                    |   |
|--------------------|---|
| Form               | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide  |
| Antibody Type      | Primary Antibody  |
| Application Dilute | Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT  |
| Application Notes  | Optimal dilution of the hCG Holo antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |