

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Thymidylate Synthase Antibody Cocktail, Clone: [TS106 + TMS715], Mouse, Monoclonal NSJ-V3164IHC-7ML

Article Name	Thymidylate Synthase Antibody Cocktail, Clone: [TS106 + TMS715], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V3164IHC-7ML
Supplier Catalog Number	V3164IHC-7ML
Alternative Catalog Number	NSJ-V3164IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogens for this Thymidylate Synthase antibody cocktail.
Product Description	It recognizes a protein of 36kDa, identified as Thymidylate Synthase (TS) (EC 2.1.1.45). TS converts deoxyuridine monophosphate (dUMP) to deoxythymidine monophosphate (dTMP), which is essential for DNA biosynthesis. TS is also a critical target for t...
Clonality	Monoclonal
Clone Designation	[TS106 + TMS715]
UniProt	P04818
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The optimal dilution of the Thymidylate Synthase antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.