

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### ATRX Antibody / RAD54, Clone: [39f], Mouse, Monoclonal NSJ-V3738SAF-100UG

|                            |   |
|----------------------------|---|
| Article Name               | ATRX Antibody / RAD54, Clone: [39f], Mouse, Monoclonal  |
| Biozol Catalog Number      | NSJ-V3738SAF-100UG  |
| Supplier Catalog Number    | V3738SAF-100UG  |
| Alternative Catalog Number | NSJ-V3738SAF-100UG  |
| Manufacturer               | NSJ Bioreagents   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC-P   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant human ATRX protein was used as the immunogen for this ATRX antibody.  |
| Product Description        | ATRX is a member of the Snf2 family of helicase/ATPases, which contribute to the remodeling of the nucleosome structure in an ATP-dependent manner, and facilitate the initiation of transcription and replication. Structurally, it contains a PHD zinc f... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [39f]   |
| UniProt                    | <a href="#">P46100</a>  |
| Purity                     | Protein G affinity chromatography   |
| Form                       | 1 mg/ml in 1X PBS, BSA free, sodium azide free  |

|                    |   |
|--------------------|---|
| Antibody Type      | Primary Antibody  |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT  |
| Application Notes  | <p>The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the ATRX antibody to be titered up or down for optimal performance.</p> <p>1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |