

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### CD74 Antibody, Clone: [CDLA74-1], Mouse, Monoclonal NSJ-V7191IHC-7ML

|                            |   |
|----------------------------|---|
| Article Name               | CD74 Antibody, Clone: [CDLA74-1], Mouse, Monoclonal   |
| Biozol Catalog Number      | NSJ-V7191IHC-7ML  |
| Supplier Catalog Number    | V7191IHC-7ML  |
| Alternative Catalog Number | NSJ-V7191IHC-7ML  |
| Manufacturer               | NSJ Bioreagents   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC   |
| Immunogen                  | Recombinant full-length human protein was used as the immunogen for the CD74 antibody. The epitope is known to be extracellular.  |
| Product Description        | CD74 associates with class II major histocompatibility complex (MHC) and is an important chaperone that regulates antigen presentation for immune response. It also serves as cell surface receptor for the cytokine macrophage migration inhibitory facto... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [CDLA74-1]  |
| UniProt                    | <a href="#">P04233</a>  |
| Purity                     | Protein G affinity chromatography   |
| Form                       | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*  |

|                    |  |
|--------------------|--|
| Antibody Type      | Primary Antibody   |
| Application Dilute | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.  |
| Application Notes  | Titering of the CD74 antibody may be required for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |