

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### RORC Antibody / ROR gamma, Clone: [RORC/2941], Mouse, Monoclonal NSJ-V7575-100UG

Article Name	RORC Antibody / ROR gamma, Clone: [RORC/2941], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V7575-100UG
Supplier Catalog Number	V7575-100UG
Alternative Catalog Number	NSJ-V7575-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P
Species Reactivity	Human
Immunogen	Recombinant human RORC protein was used as the immunogen for the RORC antibody. The epitope has been found to be with amino acids 1-50.
Product Description	This MAb recognizes a protein of 63kDa, identified as ROR-C. Its epitope maps in between aa1-50. The nuclear orphan receptors ROR alpha and ROR gamma are members of the nuclear hormone receptor superfamily. Members of this family act by directly asso...
Clonality	Monoclonal
Clone Designation	[RORC/2941]
UniProt	<a href="#">P51449</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 1-2ug/million cells,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT,Prediluted IHC only format: incubate for 30 min at RT (1)
Application Notes	Optimal dilution of the RORC antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.