

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MALT1 Antibody, Clone: [SPM578], Mouse, Monoclonal NSJ-V9006-20UG

Article Name	MALT1 Antibody, Clone: [SPM578], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9006-20UG
Supplier Catalog Number	V9006-20UG
Alternative Catalog Number	NSJ-V9006-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Amino acids 701-808 from the human protein were used as the immunogen for this MALT1 antibody.
Product Description	Mucosa associated lymphoid tissue lymphoma translocation gene 1 (MALT-1) is found in extranodal low-grade B cell lymphomas. The gene encodes two Ig-like C2-type domains and fuses with an API2 gene, which is highly expressed in adult lymphoid tissue. ...
Clonality	Monoclonal
Clone Designation	[SPM578]
UniProt	Q9UDY8
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	<p>The optimal dilution of the MALT1 antibody for each application should be determined by the researcher.</p> <ol style="list-style-type: none"> 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.