

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD35 Antibody (Complement Receptor 1), Clone: [SPM554], Mouse, Monoclonal NSJ-V9010IHC-7ML**

Article Name	CD35 Antibody (Complement Receptor 1), Clone: [SPM554], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9010IHC-7ML
Supplier Catalog Number	V9010IHC-7ML
Alternative Catalog Number	NSJ-V9010IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Human monocytes were used as the immunogen for this CD35 antibody.
Product Description	Recognizes a protein of 210-220kDa, which is identified as the Complement receptor 1 or CD35. This mAb is specific for a site in CR1 that is distal from the C3b-binding site, so that it is unable to block CR1/CD35 activity. It is highly specific and ...
Clonality	Monoclonal
Clone Designation	[SPM554]
UniProt	<a href="#">P17927</a>
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The optimal dilution of the CD35 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.