

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 8 Antibody, Clone: [SPM538], Mouse, Monoclonal NSJ-V9038-100UG

Article Name	Cytokeratin 8 Antibody, Clone: [SPM538], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9038-100UG
Supplier Catalog Number	V9038-100UG
Alternative Catalog Number	NSJ-V9038-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P
Species Reactivity	Human
Immunogen	A cytoskeleton preparation containing keratin 8 was used as the immunogen for this Cytokeratin 8 antibody.
Product Description	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc...
Clonality	Monoclonal
Clone Designation	[SPM538]
UniProt	P05787
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 1-2ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	<p>The optimal dilution of the Cytokeratin 8 antibody for each application should be determined by the researcher.</p> <ol style="list-style-type: none"> 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.