

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mucin-5AC Antibody, Clone: [SPM297], Mouse, Monoclonal NSJ-V9051SAF-100UG

Article Name	Mucin-5AC Antibody, Clone: [SPM297], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9051SAF-100UG
Supplier Catalog Number	V9051SAF-100UG
Alternative Catalog Number	NSJ-V9051SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	An M1 mucin preparation from the fluid of an ovarian mucinous cyst belonging to an O Le(a-b) patient was used as the immunogen for this Mucin-5AC antibody.
Product Description	This mAb recognizes the peptide core of gastric mucin M1 (>1,000kDa) (recently identified as Mucin 5AC). Its epitope is destroyed by beta-mercaptoethanol and proteases but not by periodate treatment. Antibody to gastric mucin M1 reacts with the gastr...
Clonality	Monoclonal
Clone Designation	[SPM297]
UniProt	P98088
Purity	Protein G affinity chromatography

Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	<p>The optimal dilution of the Mucin-5AC antibody for each application should be determined by the researcher.</p> <ol style="list-style-type: none"> 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.