

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

NKX2.2 Antibody, Clone: [SPM564], Mouse, Monoclonal NSJ-V9061-100UG

| | |
|----------------------------|---|
| Article Name | NKX2.2 Antibody, Clone: [SPM564], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V9061-100UG |
| Supplier Catalog Number | V9061-100UG |
| Alternative Catalog Number | NSJ-V9061-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC-P |
| Species Reactivity | Human |
| Immunogen | Human recombinant protein was used as the immunogen for this NKX2.2 antibody. |
| Product Description | Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s... |
| Clonality | Monoclonal |
| Clone Designation | [SPM564] |
| UniProt | O95096 |
| Purity | Protein G affinity chromatography |

| | |
|--------------------|--|
| Form | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2) |
| Application Notes | <p>The optimal dilution of the NKX2.2 antibody for each application should be determined by the researcher.</p> <ol style="list-style-type: none"> 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |