

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-TG Antibody, Clone: [SPM517], Mouse, Monoclonal NSJ-V9084-20UG

Article Name	Anti-TG Antibody, Clone: [SPM517], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9084-20UG
Supplier Catalog Number	V9084-20UG
Alternative Catalog Number	NSJ-V9084-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Human thyroid follicular cells were used as the immunogen for this anti-TG antibody.
Product Description	Thyroglobulin (TG) is a 660kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers under...
Clonality	Monoclonal
Clone Designation	[SPM517]
UniProt	<a href="#">P01266</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	The optimal dilution of the anti-TG antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.