

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-p53 Antibody, Clone: [SPM590], Mouse, Monoclonal NSJ-V9090-100UG

| | |
|----------------------------|---|
| Article Name | Anti-p53 Antibody, Clone: [SPM590], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V9090-100UG |
| Supplier Catalog Number | V9090-100UG |
| Alternative Catalog Number | NSJ-V9090-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC-P, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant human wild type p53 protein was used as the immunogen for this anti-p53 antibody. Its epitope maps within the N-terminus (approx. amino acids 37-45). |
| Product Description | Recognizes a 53kDa protein, which is identified as p53 suppressor gene product. It reacts with the mutant as well as the wild form of p53. p53 is a tumor suppressor gene expressed in a wide variety of tissue types and is involved in regulating cell g... |
| Clonality | Monoclonal |
| Clone Designation | [SPM590] |
| UniProt | P04637 |
| Purity | Protein G affinity chromatography |

| | |
|--------------------|--|
| Form | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Dilute | Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Application Notes | The optimal dilution of the anti-p53 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |