

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Tyrosinase Antibody, Clone: [SPM360], Mouse, Monoclonal NSJ-V9093IHC-7ML

Article Name	Tyrosinase Antibody, Clone: [SPM360], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9093IHC-7ML
Supplier Catalog Number	V9093IHC-7ML
Alternative Catalog Number	NSJ-V9093IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant tyrosinase protein was used as the immunogen for this Tyrosinase antibody.
Product Description	Recognizes a cluster of proteins between 70-80kDa, identified as tyrosinase. Occasionally a minor band at 55kDa is also detected. This mAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-contai...
Clonality	Monoclonal
Clone Designation	[SPM360]
UniProt	P14679
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The optimal dilution of the Tyrosinase antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, or 10mM Citrate buffer, pH6.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.