

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-SUMO-1 Antibody, Clone: [SPM571], Mouse, Monoclonal NSJ-V9095-20UG

Article Name	Anti-SUMO-1 Antibody, Clone: [SPM571], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9095-20UG
Supplier Catalog Number	V9095-20UG
Alternative Catalog Number	NSJ-V9095-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for this anti-SUMO-1 antibody.
Product Description	This mAb is specific to SUMO-1 and shows no cross-reaction with either SUMO2 or SUMO3. The small ubiquitin-related modifier (SUMO) proteins, which include SUMO1, SUMO2 and SUMO3, belong to the ubiquitin-like protein family. Like ubiquitin, the SUMO p...
Clonality	Monoclonal
Clone Designation	[SPM571]
UniProt	<a href="#">P63165</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	<p>The optimal dilution of the anti-SUMO-1 antibody for each application should be determined by the researcher.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>