

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-PGP9.5 Antibody, Clone: [SPM574], Mouse, Monoclonal NSJ-V9096-20UG

Article Name	Anti-PGP9.5 Antibody, Clone: [SPM574], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9096-20UG
Supplier Catalog Number	V9096-20UG
Alternative Catalog Number	NSJ-V9096-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IF, IHC-P, WB
Species Reactivity	Human, Rat
Immunogen	Native UchL1/PGP9.5 protein isolated from human brain was used as the immunogen for this anti-PGP9.5 antibody.
Product Description	This mAb reacts with a protein of 20-30kDa, identified as PGP9.5, also known as ubiquitin carboxyl-terminal hydrolase-1 (UchL1). Initially, PGP9.5 expression in normal tissues was reported in neurons and neuroendocrine cells but later it was found in...
Clonality	Monoclonal
Clone Designation	[SPM574]
UniProt	P09936
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Western blot: 1-2ug/ml,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	The optimal dilution of the anti-PGP9.5 antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.