

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Carbonic Anhydrase IX Antibody, Clone: [SPM487], Mouse, Monoclonal NSJ-V9101IHC-7ML

Article Name	Carbonic Anhydrase IX Antibody, Clone: [SPM487], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9101IHC-7ML
Supplier Catalog Number	V9101IHC-7ML
Alternative Catalog Number	NSJ-V9101IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	The microsomal fraction of human renal cortical tissue homogenate was used as the immunogen for this Carbonic Anhydrase IX antibody.
Product Description	Recognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/CA9/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis bloo...
Clonality	Monoclonal
Clone Designation	[SPM487]
UniProt	Q16790

Purity	Protein G affinity chromatography
Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The optimal dilution of the Carbonic Anhydrase IX antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.