

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Bcl10 Antibody, Clone: [SPM520], Mouse, Monoclonal NSJ-V9102SAF-100UG

|                            |   |
|----------------------------|---|
| Article Name               | Bcl10 Antibody, Clone: [SPM520], Mouse, Monoclonal  |
| Biozol Catalog Number      | NSJ-V9102SAF-100UG  |
| Supplier Catalog Number    | V9102SAF-100UG  |
| Alternative Catalog Number | NSJ-V9102SAF-100UG  |
| Manufacturer               | NSJ Bioreagents   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC-P   |
| Species Reactivity         | Human   |
| Immunogen                  | Amino acids 122-168 were used as the immunogen for this Bcl10 antibody.   |
| Product Description        | Bcl10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of Bcl10... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [SPM520]  |
| UniProt                    | <a href="#">O95999</a>  |
| Purity                     | Protein G affinity chromatography   |
| Form                       | 1 mg/ml in 1X PBS, BSA free, sodium azide free  |

|                    |  |
|--------------------|--|
| Antibody Type      | Primary Antibody   |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2)   |
| Application Notes  | <p>The optimal dilution of the Bcl10 antibody for each application should be determined by the researcher.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |