

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CD5 Antibody, Clone: [SPM546], Mouse, Monoclonal NSJ-V9103IHC-7ML

| | |
|----------------------------|---|
| Article Name | Anti-CD5 Antibody, Clone: [SPM546], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V9103IHC-7ML |
| Supplier Catalog Number | V9103IHC-7ML |
| Alternative Catalog Number | NSJ-V9103IHC-7ML |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC-P |
| Species Reactivity | Human |
| Immunogen | Human CD5 recombinant protein was used as the immunogen for this anti-CD5 antibody. |
| Product Description | Recognizes a 67kDa transmembrane protein which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas. Anti-CD5 is a pan T-cell mark... |
| Clonality | Monoclonal |
| Clone Designation | [SPM546] |
| UniProt | P06127 |
| Purity | Protein G affinity chromatography |

| | |
|--------------------|--|
| Form | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Dilute | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Application Notes | The optimal dilution of the anti-CD5 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |