

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CD79a Antibody, Clone: [SPM550], Mouse, Monoclonal NSJ-V9110IHC-7ML

Article Name	Anti-CD79a Antibody, Clone: [SPM550], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9110IHC-7ML
Supplier Catalog Number	V9110IHC-7ML
Alternative Catalog Number	NSJ-V9110IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Amino acids 202-216 (GTYQDVGSNIADVQ) of the human protein was used as the immunogen for this anti-CD79a antibody.
Product Description	A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B...
Clonality	Monoclonal
Clone Designation	[SPM550]
UniProt	P11912
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The optimal dilution of the anti-CD79a antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.