

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Golgi Antibody, Clone: [SPM581], Mouse, Monoclonal NSJ-V9111-100UG

Article Name	Anti-Golgi Antibody, Clone: [SPM581], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9111-100UG
Supplier Catalog Number	V9111-100UG
Alternative Catalog Number	NSJ-V9111-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	SU-DHL-1 large cell lymphoma cells were used as the immunogen for this anti-Golgi antibody.
Product Description	This mAb recognizes Golgi complex in human cells. It is a part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, an...
Clonality	Monoclonal
Clone Designation	[SPM581]
UniProt	Not Known
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	<p>The optimal dilution of the anti-Golgi antibody for each application should be determined by the researcher.</p> <ol style="list-style-type: none"> 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.