

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Hepatocyte Antibody, Clone: [SPM582], Mouse, Monoclonal NSJ-V9119IHC-7ML

Article Name	Hepatocyte Antibody, Clone: [SPM582], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9119IHC-7ML
Supplier Catalog Number	V9119IHC-7ML
Alternative Catalog Number	NSJ-V9119IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Extract of a formalin-fixed rejected-allograft of human liver was used as the immunogen for this Hepatocyte antibody.
Product Description	Hepatocyte Specific Antigen, also called Hepatocyte Paraffin 1 or Hep Par 1, localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholang...
Clonality	Monoclonal
Clone Designation	[SPM582]
UniProt	Not Known
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The optimal dilution of the Hepatocyte antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.