

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **NOC4L Antibody, Clone: [PCRP-NOC4L-1E3], Mouse, Monoclonal NSJ-V9212-20UG**

|                            |   |
|----------------------------|---|
| Article Name               | NOC4L Antibody, Clone: [PCRP-NOC4L-1E3], Mouse, Monoclonal  |
| Biozol Catalog Number      | NSJ-V9212-20UG  |
| Supplier Catalog Number    | V9212-20UG  |
| Alternative Catalog Number | NSJ-V9212-20UG  |
| Manufacturer               | NSJ Bioreagents   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FACS, IF  |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant full-length human NOC4L protein was used as the immunogen for the NOC4L antibody.   |
| Product Description        | Treg activation is critical for maintaining self-tolerance, but the translational control of this process is still poorly understood. Zhu et al. report a conserved ribosome biogenesis factor, Noc4L, that regulates mRNAs related to Treg and Tconv acti... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [PCRP-NOC4L-1E3]  |
| UniProt                    | <a href="#">Q9BVI4</a>  |
| Purity                     | Protein A/G affinity  |

|                    |  |
|--------------------|--|
| Form               | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide        |
| Antibody Type      | Primary Antibody   |
| Application Dilute | Flow cytometry: 1-2ug/million cells,Immunofluorescence: 1-2ug/ml               |
| Application Notes  | Optimal dilution of the NOC4L antibody should be determined by the researcher. |