

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Dog sRANKL(Soluble Receptor Activator of Nuclear factor-**κ**B Ligand) ELISA Kit EBT-ELK9165

Article Name	Dog sRANKL(Soluble Receptor Activator of Nuclear factor- κ B Ligand) ELISA Kit
Biozol Catalog Number	EBT-ELK9165
Supplier Catalog Number	ELK9165
Alternative Catalog Number	EBT-ELK9165-96, EBT-ELK9165-48, EBT-ELK9165-96X5
Manufacturer	ELK Biotechnology
Category	Kits/Assays
Species Reactivity	Canine
Concentration	1000 pg/mL
Range	15.63-1000 pg/mL
Sensitivity	5.5 pg/mL
Samples	serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids

Application Notes

Assay Type: Sandwich. Assay length: 3.5h. Research Area: Signal transduction, Metabolic pathway, Apoptosis, Tumor immunity, Bone metabolism,. Test principle: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Dog sRANKL. Standards or samples are added to the appropriate microtiter plate wells then with a biotin-conjugated antibody specific to Dog sRANKL. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Dog sRANKL, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm 10nm. The concentration of Dog sRANKL in the samples is then determined by comparing the OD of the samples to the standard curve