

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Mouse CD47 (C-Fc) EBT-EPT116

Article Name	Recombinant Mouse CD47 (C-Fc)
Biozol Catalog Number	EBT-EPT116
Supplier Catalog Number	EPT116
Alternative Catalog Number	EBT-EPT116-10
Manufacturer	ELK Biotechnology
Category	Proteine/Peptide
Product Description	Recombinant Mouse Leukocyte Surface Antigen CD47 is produced by our Mammalian expression system and the target gene encoding Gln19-Pro158 is expressed with a Fc tag at the C-terminus....
Molecular Weight	Molecular weight: 42.8 KDa. Apparent molecular weight: 60-90 KDa, reducing conditions
UniProt	Q61735
Purity	Greater than 95% as determined by reducing SDS-PAGE.

Application Notes

Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test. Background: CD47, also known as Integrin-Associated Protein (IAP) and OA3, is a glycosylated atypical member of the immunoglobulin superfamily. Mouse CD47 is an integral membrane protein that consists of a extracellular domain (ECD) with a single Ig-like domain, five membrane-spanning regions with short intervening loops, and C-terminal cytoplasmic tail. CD47 has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. It plays an important role in memory formation and synaptic plasticity in the hippocampus. As a receptor for SIRPA, its binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Interaction with SIRPG mediates cell-cell adhesion, it enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation. It may play a role in membrane transport and/or integrin dependent signal transduction. It also prevents premature elimination of red blood cells