

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Mouse IL-22 (C-mFc) EBT-EPT148

Article Name	Recombinant Mouse IL-22 (C-mFc)
Biozol Catalog Number	EBT-EPT148
Supplier Catalog Number	EPT148
Alternative Catalog Number	EBT-EPT148-10
Manufacturer	ELK Biotechnology
Category	Proteine/Peptide
Product Description	Recombinant Mouse Interleukin-22 is produced by our Mammalian expression system and the target gene encoding Leu34-Val179 is expressed with a mFc tag at the C-terminus....
Molecular Weight	Molecular weight: 43.8 KDa. Apparent molecular weight: 50-65 KDa, reducing conditions
UniProt	Q9JJY9
Purity	Greater than 95% as determined by reducing SDS-PAGE.

Application Notes

Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL test. Background: Interleukin-22 (IL-22) was initially identified as a gene induced by IL-9 in mouse T cells and mast cells. Mouse IL-22 cDNA encodes a 179 amino acid residue protein with a putative 33 amino acid signal peptide that is cleaved to generate a 147 amino acid mature protein that shares approximately 79% and 22% sequence identity with human IL22 and IL10, respectively. IL22 has been shown to activate STAT-1 and STAT-3 in several hepatoma cell lines and up-regulate the production of acute phase proteins. IL-22 is produced by normal mouse T cells upon Con A activation. Mouse IL-22 expression is also induced in various organs upon lipopolysaccharide injection, suggesting that IL-22 may be involved in inflammatory responses. The functional IL-22 receptor complex consists of two receptor subunits, IL-22R (previously an orphan receptor named CRF2-9) and IL-10R β (previously known as CRF2-4), belonging to the class II cytokine receptor family