

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Human APRIL (N-Flag-His) EBT-EPT169

Article Name	Recombinant Human APRIL (N-Flag-His)
Biozol Catalog Number	EBT-EPT169
Supplier Catalog Number	EPT169
Alternative Catalog Number	EBT-EPT169-10
Manufacturer	ELK Biotechnology
Category	Proteine/Peptide
Product Description	Recombinant Human A proliferation-inducing Ligand is produced by our Mammalian expression system and the target gene encoding Lys112-Leu250 is expressed with a His, Flag tag at the N-terminus....
Molecular Weight	Molecular weight: 50 KDa. Apparent molecular weight: 60 KDa, reducing conditions
UniProt	O75888
Purity	Greater than 95% as determined by reducing SDS-PAGE.

Application Notes

Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL test. Biological activity: Immobilized Human APRIL -Flag-His(CatCU89) at 1 μ g/ml (100 μ l/well) can bind Cynomolgus BCMA-Fc(CatCP59).The ED50 of Cynomolgus BCMA-Fc(CatCP59) is 4.23 ng/ml. Background: APRIL (a proliferation-inducing ligand), also known as TNFSF13, TALL2, TRDL1, and CD256, is a member of the TNF ligand superfamily. It is synthesized as a 32 kDa proprotein which is cleaved by furin in the Golgi to release the active 17 kDa soluble molecule. Secreted human APRIL, which consists almost entirely of a single TNF homology domain, shares 85% amino acid sequence identity with mouse and rat APRIL. Both APRIL and its close relative BAFF bind and signal through the TNF superfamily receptors TACI and BCMA, while BAFF additionally functions through BAFF R. APRIL binds to heparan sulfate proteoglycans (HSPGs) independently of its binding to TACI and BCMA. APRIL can form bioactive heterotrimers with BAFF, and these circulate in the serum of patients with rheumatic immune disorders. APRIL enhances the proliferation and survival of plasma cells and also promotes T cell-dependent humoral responses. APRIL levels are elevated in the serum during coronary artery disease, and it is also elevated in many cancers primarily due to expression by tumor-infiltrating neutrophils