

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Human Siglec-5 (C-6His-Flag-Fc) EBT-EPT208

Article Name	Recombinant Human Siglec-5 (C-6His-Flag-Fc)
Biozol Catalog Number	EBT-EPT208
Supplier Catalog Number	EPT208
Alternative Catalog Number	EBT-EPT208-10
Manufacturer	ELK Biotechnology
Category	Proteine/Peptide
Product Description	Recombinant Human Sialic Acid-binding Ig-like Lectin 5 is produced by our Mammalian expression system and the target gene encoding Glu17-Thr434 is expressed with a 6His, Flag, Fc tag at the C-terminus....
Molecular Weight	Molecular weight: 74.1 KDa. Apparent molecular weight: 90-110 KDa, reducing conditions
UniProt	O15389
Purity	Greater than 95% as determined by reducing SDS-PAGE.

Application Notes

Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL test. Background: Human Siglec-5 are Igtype(Igtype) lectins belonging to the Ig superfamily, They are characterized by an N terminal Ig-like V type domain which mediates sialic acid binding, followed by varying numbers of Ig-like C2 type domains. SIGLEC5 has also been designated CD170, they are expressed by monocytic or myeloid lineage cells, and also found at high levels in peripheral blood leukocytes, spleen, bone marrow and at lower levels in lymph node, lung, appendix, placenta, pancreas and thymus. SIGLEC5 are expressed by monocytes and neutrophils but absent from leukemic cell lines representing early stages of myelomonocytic differentiation. Siglec5 to 11 share a high degree of sequence similarity with CD33/Siglec3 both in their extracellular and intracellular regions. They are collectively referred to as CD33 related Siglecs. One remarkable feature of the CD33 related Siglecs is their differential expression pattern within the hematopoietic system. This fact, together with the presence of two conserved immunoreceptor tyrosinebased inhibition motifs (ITIMs) in their cytoplasma tails, suggests that CD33 related Siglecs are involved in the regulation of cellular activation within the immune system