

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Recombinant Rabbit Monoclonal Antibody to TNF-alpha (Tumor Necrosis Factor alpha)(Clone : TNF/1500R), Clone: [TNF/1500R] ABI-12-1042

|                            |   |
|----------------------------|---|
| Article Name               | Recombinant Rabbit Monoclonal Antibody to TNF-alpha (Tumor Necrosis Factor alpha)(Clone : TNF/1500R), Clone: [TNF/1500R]  |
| Biozol Catalog Number      | ABI-12-1042   |
| Supplier Catalog Number    | 12-1042   |
| Alternative Catalog Number | ABI-12-1042-100UG   |
| Manufacturer               | Abeomics  |
| Category                   | Sonstiges   |
| Application                | IHC   |
| Species Reactivity         | Human, Rat  |
| Immunogen                  | Recombinant full-length human TNF-alpha protein   |
| Product Description        | This MAb recognizes human 17-26kDa protein, which is identified as cytokine TNF-alpha (Tumor Necrosis Factor-alpha). TNF-alpha can be expressed as a 17kDa free molecule, or as a 26kDa membrane protein. TNF-alpha is a protein secreted by lipopolysacch... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [TNF/1500R]   |
| NCBI                       | <a href="#">7124</a>  |
| UniProt                    | <a href="#">P01375</a>  |
| Purity                     | Purified Ab with BSA and Azide at 200ug/ml  |

|                    |  |
|--------------------|--|
| Form               | Purified   |
| Target             | TNF-alpha (Tumor Necrosis Factor alpha)  |
| Antibody Type      | Recombinant Antibody   |
| Application Dilute | Immunohistochemistry (Formalin-fixed) (4-8µg/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes), |